

On well quasiordering of finite languages

Martin Klazar*

Department of Applied Mathematics of Charles University, Malostranské náměstí 25, 11800 Praha 1,
Czech Republic

Received 8 July 1993; revised 29 August 1995

Abstract

We investigate here the quasiordering \preceq of finite sets of finite strings over an infinite set of symbols S . We set $\mathcal{X} \preceq \mathcal{L}$ iff it is possible to rename symbols occurring in the strings of \mathcal{L} so that any string of \mathcal{X} is a subsequence of a string of the renamed \mathcal{L} . We prove that \preceq is a wqo which answers the question raised by Gustedt (1992). We prove also a stronger version with injective correspondence between strings.

1. Introduction

Strings are finite sequences over S where S is an infinite countable set of *symbols*. *Languages* are finite sets of strings, *babels* are sets of languages. If $A \subseteq S$ then A^* stands for the set of all strings over A . By S^{**} we denote the babel consisting of all languages. We define, for a string $u = a_0 a_1 \dots a_m$, $S(u) = \bigcup_{i=0}^m \{a_i\}$. Similarly, $S(\mathcal{L}) = \bigcup_{u \in \mathcal{L}} S(u)$ for any language \mathcal{L} .

The notation $u \subset v$ means, for any two sequences $u = a_0 a_1 \dots a_m$ and $v = b_0 b_1 \dots b_n$, that u is a subsequence of v : $a_0 = b_{j_0}$, $a_1 = b_{j_1}$, \dots , $a_m = b_{j_m}$ for some m indices $0 \leq j_0 < j_1 < \dots < j_m \leq n$. We define, for two languages \mathcal{L} and \mathcal{X} , that $\mathcal{L} \preceq \mathcal{X}$ (via f) iff $u \subset f(u)$ for any $u \in \mathcal{L}$ for some mapping $f: \mathcal{L} \rightarrow \mathcal{X}$. A mapping $\varphi: S \rightarrow S$ transforms a language \mathcal{L} to the language $\varphi(\mathcal{L}) = \{\varphi(u) \mid u \in \mathcal{L}\}$ where $\varphi(u) = \varphi(a_0 a_1 \dots a_m) = \varphi(a_0) \varphi(a_1) \dots \varphi(a_m)$. We shall investigate the following quasiordering.

Definition 1.1. $\mathcal{L} \preceq \mathcal{X}$, \mathcal{L} and \mathcal{X} are languages, iff $\mathcal{L} \preceq \varphi(\mathcal{X})$ for some $\varphi: S \rightarrow S$.

The above quasiordering was introduced in [5] to generalize *chain minor* ordering of finite posets. We say, in accordance with [5] and with [2] that P is a chain minor of

* E-mail: klazar@kam.ms.mff.cuni.cz.

Q (P and Q are finite posets) iff there is a mapping $\rho: Q \rightarrow P$ such that any chain in P is isomorphic via ρ to a chain in Q (thus ρ must be onto). Chain minor ordering was introduced in connection with scheduling stochastic project networks [5]. Clearly, P is a chain minor of Q iff $\mathcal{L}(P) \leq \mathcal{L}(Q)$ where $\mathcal{L}(P)$ and $\mathcal{L}(Q)$ are languages consisting of chains in corresponding posets.

By means of that equivalence it has been proven in [2] (see also [1]) that chain minor is a wqo of finite posets. The proof uses substantially the fact that any ‘poset language’ $\mathcal{L}(P)$ consists of strings without repetitions. The problem whether \leq is a wqo for languages in general was posed [2]. Generalizing the approach in [2] we answer this question affirmatively.

Theorem 1.2. (S^{**}, \leq) is a wqo.

One can define a stronger quasiordering \leq^* if the mapping f in the definition of \leq is injective in addition. We prove that \leq^* is a wqo as well.

Theorem 1.3. (S^{**}, \leq^*) is a wqo.

In Section 2 we give some preliminaries and demonstrate in a simple case our method. Theorems 1.2 and 1.3 are proved in Sections 3 and 4, respectively. In Section 5 we give counterexamples showing that requiring an injective ϕ in Definition 1.1 destroys the wqo property.

2. Absolute minimum about wqo

Any transitive and reflexive binary relation is called a *quasiordering* or, shortly, *qo*. If (Q, \leq_Q) is a qo then $x <_Q y$ means that $x \leq_Q y$ and $y \not\leq_Q x$. A cone determined by the element $x \in Q$ is the set $K_x = \{y \in Q \mid y \geq_Q x\}$. A qo (Q, \leq_Q) is a *well quasiordering* or, shortly, *wqo* if it possesses the property characterized by the following lemma. For the proof and for more background we refer to [4].

Lemma 2.1. Suppose (Q, \leq_Q) is a qo. The following conditions are equivalent.

- (1) For any infinite sequence $(q_i)_{i=0}^\infty \subseteq Q$ there are indices $i < j$ such that $q_i \leq_Q q_j$.
- (2) For any infinite sequence $(q_i)_{i=0}^\infty \subseteq Q$ there are indices $0 \leq i_0 < i_1 < \dots$ such that

$$q_{i_0} \leq_Q q_{i_1} \leq_Q \dots$$

- (3) No infinitely many elements x_0, x_1, \dots of Q create an antichain or a strictly descending chain

$$x_0 >_Q x_1 >_Q \dots$$

Sequences satisfying (1) are called *good*, other sequences are called *bad*. The infinite monotonic subsequence in (2) is called *perfect*. We recall two folcloric but useful statements.

Cone deleting argument. Suppose (Q, \leq_Q) is a wqo and Q_0, Q_1, \dots are defined by $Q_0 = Q, Q_{i+1} = Q_i \setminus K_{q_i}, q_i \in Q_i$. Then this sequence is finite, $Q_j = \emptyset$ for some j (otherwise $(q_i)_{i=0}^\infty \subseteq Q$ would be a bad sequence).

Product argument. Suppose $(Q_i, \leq_{Q_i})_{i=0}^r$ are wqo's, $Q = Q_0 \times Q_1 \times \dots \times Q_r$ and (Q, \leq_{pr}) is defined by $(x_i)_{i=0}^r \leq_{pr} (y_i)_{i=0}^r$ iff $x_i \leq_{Q_i} y_i$ for $i = 0, \dots, r$. Then (Q, \leq_{pr}) is a wqo as well (apply Lemma 2.1 $r + 1$ times).

Let (Q, \leq_Q) be a qo. The *Higman ordering* $(SEQ(Q), \leq_H)$ on the set

$$SEQ(Q) = \{(I, \ell) \mid I \text{ is a finite linear ordering and } \ell: I \rightarrow Q\}$$

of all finite sequences over Q is defined by $(I_0, \ell_0) \leq_H (I_1, \ell_1)$ iff there is an increasing mapping $F: I_0 \rightarrow I_1$ such that $\ell_0(x) \leq_Q \ell_1(F(x))$ for any $x \in I_0$. We will use the following classical result of the wqo theory [3].

Theorem 2.2 (Higman [3]). $(SEQ(Q), \leq_H)$ is a wqo for any wqo (Q, \leq_Q) .

To demonstrate our method in a simple case we prove as an example a weaker version of Higman theorem which deals with the structure $(SET(Q), \leq_S)$ consisting of finite subsets of Q with the qo $A \leq_S B$ iff there is an injective mapping $F: A \rightarrow B$ such that $x \leq_Q F(x)$ for any $x \in A$.

Lemma 2.3. $(SET(Q), \leq_S)$ is a wqo for any wqo (Q, \leq_Q) .

Proof. We prove by a direct argument that any sequence $A = (A_i)_{i=0}^\infty \subseteq SET(Q)$ is good to \leq_S . We say that $X = (B_i, C_i)_{i=0}^\infty$ is a *friend* of A if $(B_i)_{i=0}^\infty$ is a subsequence of $A, C_i \subseteq B_i$ for any i , and $(|C_i|)_{i=0}^\infty$ is bounded. Set $R(X) = \bigcup_{i=0}^\infty (B_i \setminus C_i)$ and $G(i, x) = |K_x \cap (B_i \setminus C_i)|$ where $x \in Q$. We say that X is a *good friend* of A if in addition $\lim_{i \rightarrow \infty} G(i, x) = \infty$ (i.e., for any m there is an n such that $i \geq n$ implies $G(i, x) \geq m$) for any fixed $x \in R(X)$.

To prove that any A has a good friend we define a (finite) sequence X_0, X_1, \dots of friends of A and initiate it by $X_0 = (A_i, \emptyset)_{i=0}^\infty$. Suppose that $X_k = (B_i, C_i)_{i=0}^\infty$ is a friend of A which fails to be a good friend: $G(i_0, x), G(i_1, x), \dots \leq N < \infty$ for some indices $0 \leq i_0 \leq i_1 < \dots$ and some $x \in R(X_k)$. Let $D_i = C_i \cup (K_x \cap (B_i \setminus C_i))$. Then

$$X_{k+1} = (B_i, D_i)_{i=0}^\infty$$

is a friend of A and moreover $R(X_{k+1}) \subseteq R(X_k) \setminus K_x$. According to the cone deleting argument (X_0, X_1, \dots) terminates in a good friend of A . Notice that when $(|A_i|)_{i=0}^\infty$ is bounded then the good friend of A obtained is $(A_i, A_i)_{i=0}^\infty$.

So let $X = (B_i, C_i)_{i=0}^\infty$ be a good friend of A . We may assume that $(|C_i|)_{i=0}^\infty$ is constant and that $C_0 \leq_S C_1 \leq_S \dots$ because by the product argument $(C_i)_{i=0}^\infty$ contains a perfect subsequence. Take j sufficiently large such that $G(j, x) \geq |B_0 \setminus C_0|$ for any

$x \in B_0 \setminus C_0$. As $C_0 \leq_s C_j$ and any $x \in B_0 \setminus C_0$ is majorized (in \leq_Q) by sufficiently many elements in $B_j \setminus C_j$ we conclude that $B_0 \leq_s B_j - A$ is good. \square

Recall that A^* is the set of all strings over A and that \subset here is the subsequence relation. The following result is an easy and well-known consequence of Higman theorem.

Corollary 2.4. *Let A be a finite alphabet. Then (A^*, \subset) is a wqo.*

3. Proof of Theorem 1.2

Any finite collection $G = (E, I) = (E(G), I(G)) = (\{e_i \mid i \in I\}, I)$ of finite sets is called a *set system*, elements of E are called *edges*. We permit repetition of edges and for simplicity we omit the indices of edges when possible. If $H = (F, J)$ is another set system such that $F \subseteq E$ (and $J \subseteq I$) then H is said to be a *subsystem* of G . If E consists of mutually disjoint edges then G is said to be a *disjoint system*.

The *matching number* $M(G)$ of $G = (E, I)$ is defined as the maximum number of edges in a disjoint subsystem of G . A Q -system is a couple (G, ℓ) where $\ell: E(G) \rightarrow Q$ gives to the edges of G labels from the set Q .

Suppose $A = (G_i, \ell_i)_{i=0}^\infty$ is a sequence of Q -systems where (Q, \leq_Q) is a qo. We say that

$$X = (H_i, \ell_i, H'_i)_{i=0}^\infty$$

is a *friend* of A if $(H_i, \ell_i)_{i=0}^\infty$ is a subsequence of A , H'_i is a subsystem of H_i , and $(M(H'_i))_{i=0}^\infty$ is bounded.

We define further

$$R(X) = \bigcup_{i=0}^\infty \ell_i(E(H_i) \setminus E(H'_i)) \subseteq Q \quad \text{and} \quad G(i, x) = M(H'_i(x)),$$

where $x \in Q$ and $H'_i(x)$ is a subsystem of H_i consisting of the edges

$$\{e \in E(H_i) \setminus E(H'_i) \mid \ell_i(e) \in K_x\}.$$

We say that X is a *good friend* of A if in addition

$$\lim_{i \rightarrow \infty} G(i, x) = \infty$$

for any $x \in R(X)$.

Lemma 3.1. *Any sequence $A = (G_i, \ell_i)_{i=0}^\infty$ of Q -systems labelled by a wqo (Q, \leq_Q) has a good friend X .*

Proof. We define again a sequence X_0, X_1, \dots of friends of A starting with $X_0 = (G_i, \ell_i, \emptyset)_{i=0}^\infty$ and show that it terminates in a good friend of A . Suppose $X_k = (H_i, \ell_i, H'_i)_{i=0}^\infty$ fails to be a good friend of A : $G(i_0, y), G(i_1, y), \dots \leq N < \infty$ for some indices $0 \leq i_0 < i_1 < \dots$ and some $y \in R(X_k)$. Then

$$X_{k+1} = (H_i, \ell_i, H'_i \cup H''_i(y))_{i=0}^\infty$$

is clearly a new friend of A and moreover $R(X_{k+1}) \subseteq R(X_k) \setminus K_y$. According to the cone deleting argument after finitely many steps a good friend of A arises. \square

Definition 3.2. A (k, l) -babel where k, l are positive integers is any pair (\mathcal{B}, A) satisfying

1. \mathcal{B} is a babel,
2. $A \subseteq S, |A| \leq l$,
3. $|S(u) \setminus A| \leq k$ whenever $u \in \mathcal{L}, \mathcal{L} \in \mathcal{B}$.

Definition 3.3. We denote by S_A^S , $A \subseteq S$, the set of all mappings $\varphi: S \rightarrow S$ such that $\varphi|_A = \text{id}_A$ and $\varphi^{-1}(A) = A$. For two languages \mathcal{X} and \mathcal{L} the notation $\mathcal{X} \leq_A \mathcal{L}$ means that $\mathcal{X} \leq \varphi(\mathcal{L})$ for some $\varphi \in S_A^S$.

Definition 3.4. Let $\mathcal{L} = (\mathcal{L}_i)_{i=0}^\infty \subseteq \mathcal{B}$ be a sequence of languages of a (k, l) -babel (\mathcal{B}, A) . Let R be a set of k symbols disjoint to A and let $\chi \in S_A^S$ be fixed such that it maps any $S(u) \setminus A, u \in \mathcal{L}_i, i \geq 0$, injectively to R . We introduce the following sequence of Q -systems $P(\mathcal{L}) = (G_i, \ell_i)_{i=0}^\infty$ (see “Added in proof”, page 88):

$$I(G_i) = \mathcal{L}_i, \quad E(G_i) = \{S(u) \setminus A \mid u \in \mathcal{L}_i\}, \quad (Q, \leq_Q) = ((R \cup A)^*, \subset),$$

$$\ell_i(e_u) = \chi(u) = \chi(a_0 a_1 \dots a_m) = \chi(a_0) \chi(a_1) \dots \chi(a_m).$$

Observation 3.5. To prove Theorem 1.2, it suffices to prove that $((\mathcal{B}, A), \leq_A)$ is a wqo for any (k, l) -babel (\mathcal{B}, A) .

Proof. If $\mathcal{L} = (\mathcal{L}_i)_{i=0}^\infty$ is a sequence of languages and $|S(u)|, u \in \mathcal{L}_i, i \geq 0$, is not universally bounded then $|S(u_0)|$, for some u_0 in some \mathcal{L}_i , is at least as big as the sum of lengths of the strings in \mathcal{L}_0 . Then it is easy to embed the whole \mathcal{L}_0 in this single string u_0 and \mathcal{L} is good. Otherwise $|S(u)| \leq c$ for all $u \in \mathcal{L}_i$ and all $i \geq 0$ and hence \mathcal{L} is a $(c, 0)$ -babel. \square

Lemma 3.6. $((\mathcal{B}, A), \leq_A)$ is a wqo for any (k, l) -babel (\mathcal{B}, A) .

Proof. We proceed by double induction on k and l and start with $k = 0$. Then $((\mathcal{B}, A), \leq_A)$ is a wqo because even $(\text{SET}(A^*), \leq_S)$ is a wqo by Lemma 2.2 and Corollary 2.4.

Suppose now that (\mathcal{B}, A) is a (k, l) -babel, $k > 0$, and $\mathcal{L} = (\mathcal{L}_i)_{i=0}^\infty \subseteq \mathcal{B}$ is a sequence of languages. We prove that \mathcal{L} is good. We may suppose, renaming appropriately symbols, that $S(\mathcal{L}_i)$ are mutually disjoint up to A and that $S \setminus \bigcup_{i \geq 0} S(\mathcal{L}_i)$ is infinite.

Let $P(\mathcal{L}) = (G_i, \ell_i)_{i=0}^\infty$ be the sequence defined in Definition 3.4. The labels form a wqo by Corollary 2.4. Thus, there is, by Lemma 3.1, a good friend $(H_i, \ell_i, H'_i)_{i=0}^\infty$ of $P(\mathcal{L})$.

Let F_i be a maximum disjoint subsystem of H'_i and let $U_i = \bigcup E(F_i)$. Clearly, $|U_i| \leq ck$ for some constant c (the bound on matching numbers) for any $i \geq 0$. We introduce a set T , $|T| = ck$, of completely new symbols which is disjoint to A and to all $\bigcup E(H_i)$. Let $\rho \in S_A^S$ be such that ρ is an identity on $S \setminus \bigcup_{i \geq 0} U_i$ and maps any U_i injectively to T .

Consider now the babel $\mathcal{C} = (\mathcal{H}_i)_{i=0}^\infty$ where $(\mathcal{H}_i)_{i=0}^\infty$ is defined by $\mathcal{H}_i = I(H'_i)$. We see that, crucially, $(\mathcal{C}, T \cup A)$ is a $(k-1, ck+1)$ -babel because any edge of H'_i must intersect U_i . We may suppose, according to the induction hypothesis, that $\rho(\mathcal{H}_0) \leq_{A \cup T} \rho(\mathcal{H}_1) \leq_{A \cup T} \dots$.

We compare the first term with the others: there are mappings $\varphi_i \in S_{A \cup T}^S$ and $f_i: \mathcal{H}_0 \rightarrow \mathcal{H}_i$, $i \geq 1$, such that $\rho(u) \subset \varphi_i(\rho(f_i(u)))$ for any $u \in \mathcal{H}_0$. Let j be such a large number that there are $|E(H_0) \setminus E(H'_0)|$ mutually disjoint edges

$$F = \{h_e | e \in E(H_0) \setminus E(H'_0)\} \subseteq E(H_j) \setminus (H'_j)$$

satisfying $\ell_j(h_e) \supset \ell_0(e)$ for any $e \in E(H_0) \setminus E(H'_0)$ and moreover any edge of F is disjoint to $S(f_j(\mathcal{H}_0))$.

We take a mapping $\varphi \in S_A^S$ as follows.

- If $x \in S(f_j(\mathcal{H}_0)) \cap U_j$ then $\rho(y) = \rho(x)$ for at most one $y \in U_0$. If it exists we put $\varphi(x) = y$.
- If $x \in S(f_j(\mathcal{H}_0)) \setminus U_j$ then we put $\varphi(x) = \varphi_j(x)$.
- If $x \in h_e$ for $e \in E(H_0) \setminus E(H'_0)$ then $\chi(y) = \chi(x)$ for at most one $y \in e$. If it exists we put $\varphi(x) = y$.

Otherwise φ is defined arbitrarily. Clearly, $I(H_0) \leq \varphi(I(H_j))$ and we conclude that the sequence \mathcal{L} is good. \square

Lemma 3.6 and Observation 3.5 prove Theorem 1.2.

4. Proof of Theorem 1.3

An easy check shows that only in Observation 3.5, we used the fact that the mapping f of the definition of \leq had not to be injective. In Lemma 3.6, it has been proven actually that $((\mathcal{B}, A), \leq_A^*)$ is a wqo for any (k, l) -babel (\mathcal{B}, A) . Now we make the whole proof injective by replacing Observation 3.5 by a finer consideration.

Suppose $\mathcal{L} = (\mathcal{L}_i)_{i=0}^\infty \subseteq S^{**}$ is a sequence of languages. We say that $X = (\mathcal{X}_i, \mathcal{X}'_i)_{i=0}^\infty$ is a *friend* of \mathcal{L} if $(\mathcal{X}_i)_{i=0}^\infty$ is a subsequence of \mathcal{L} , $\mathcal{X}'_i \subseteq \mathcal{X}_i$, $(|\mathcal{X}'_i|)_{i=0}^\infty$ is constant, and $\min\{|S(u)| | u \in \mathcal{X}'_i\} \rightarrow \infty$ for $i \rightarrow \infty$. If moreover $(\max\{|S(u)| | u \in \mathcal{X}_i\})_{i=0}^\infty$ is bounded then X is said to be a *good friend* of \mathcal{L} .

Consider the following property:

- (*) For any c there are in some language \mathcal{L}_i c strings u such that for each of them $|S(u)| \geq c$

Lemma 4.1. Suppose $\mathcal{L} = (\mathcal{L}_i)_{i=0}^\infty$ is a sequence of languages not having property (*). Then \mathcal{L} has a good friend.

Proof. We define then by induction a sequence X_0, X_1, \dots of friends of \mathcal{L} starting with $X_0 = (\mathcal{L}_i, \emptyset)_{i=0}^\infty$. If $X_k = (\mathcal{X}_i, \mathcal{X}'_i)_{i=0}^\infty$ fails to be a good friend of \mathcal{L} then $|S(u_j)| \rightarrow \infty$ for $j \rightarrow \infty$ for some strings $u_i \in \mathcal{X}_i \setminus \mathcal{X}'_i$, and some indices $0 \leq i_0 < i_1 < \dots$. Then

$$X_{k+1} = (\mathcal{X}_i, \mathcal{X}'_i \cup \{u_{i_j}\})_{j=0}^\infty$$

is a new friend of \mathcal{L} . As (*) is violated the growth of $|\mathcal{X}'_i|$ cannot proceed arbitrarily long and after finitely many steps a good friend of \mathcal{L} is obtained. \square

Proof of Theorem 1.3: Suppose $\mathcal{L} = (\mathcal{L}_i)_{i=0}^\infty \subseteq S^{**}$ is a sequence of languages. If \mathcal{L} has property (*) then \mathcal{L}_0 embeds injectively in some \mathcal{L}_j . If not then consider a good friend $X = (\mathcal{X}_i, \mathcal{X}'_i)_{i=0}^\infty$ of \mathcal{L} . The sequence $(\mathcal{X}_i \setminus \mathcal{X}'_i)_{i=0}^\infty$ is a $(c, 0)$ -babel for some c and by Lemma 3.6 we may suppose it forms a perfect sequence $(\mathcal{X}_0 \setminus \mathcal{X}'_0) \leq^* (\mathcal{X}_1 \setminus \mathcal{X}'_1) \leq^* \dots$.

So there are mappings $\varphi_i: S \rightarrow S$ and injective mappings $f_i: (\mathcal{X}_0 \setminus \mathcal{X}'_0) \rightarrow (\mathcal{X}_i \setminus \mathcal{X}'_i)$, $i \geq 1$, such that $u \subset \varphi_i(f_i(u))$ for any $u \in \mathcal{X}_0 \setminus \mathcal{X}'_0$. Now we take such a large j that

$$\min_{u \in \mathcal{X}_j} |S(u)| \geq \sum_{v \in \mathcal{X}_0 \setminus \mathcal{X}'_0} |S(f_j(v))| + \sum_{v \in \mathcal{X}'_0} \text{length}(v).$$

It is easy to extend the injective covering $\mathcal{X}_0 \setminus \mathcal{X}'_0 \leq^* \mathcal{X}_j \setminus \mathcal{X}'_j$ to the injective covering $\mathcal{X}_0 \leq^* \mathcal{X}_j$. We conclude that \mathcal{L} is good. \square

5. Concluding remarks

Now we show that the fact we did not require an injective φ was crucial to obtain wqo. Let $\mathcal{X} \leq_* \mathcal{L}$, for two languages \mathcal{L} and \mathcal{X} , iff there is an injective $\varphi: S \rightarrow S$ such that $\mathcal{X} \leq \varphi(\mathcal{L})$. Consider this example.

Example 5.1. The infinite babels

$$\mathcal{A}_0 = \{\{132132\}, \{14213243\}, \{1521324354\}, \{162132435465\}, \dots\}$$

and

$$\mathcal{A}_1 = \{\{ab, bc, ca\}, \{ab, bc, cd, da\}, \{ab, bc, cd, de, ea\}, \dots\}$$

are antichains to \leq_* . Thus, \leq_* is not a wqo.

Note that both babels are antichains also in the ordering obtained by replacing in Definition 1.1, $\mathcal{X} \leq \varphi(\mathcal{L})$ by $\varphi(\mathcal{X}) \leq \mathcal{L}$.

Problem 5.2. Suppose now that a language $\mathcal{L} = u_0 u_1 \dots u_k$ is a finite sequence of strings rather than just a set and put $\mathcal{L} = u_0 u_1 \dots u_k \preceq \mathcal{K} = v_0 v_1 \dots v_l$ iff there is a mapping $\varphi: S \rightarrow S$ and an increasing injection $f: \{0, 1, \dots, k\} \rightarrow \{0, 1, \dots, l\}$ such that $u_i \subseteq \varphi(v_{f(i)})$ for all $i = 0, 1, \dots, k$. Is this \preceq still a wqo?

Acknowledgements

Author is grateful to Robin Thomas and especially to Jens Gustedt for introducing him to this problem. Comments of one anonymous referee helped to improve readability of the paper.

Added in proof. In Definition 3.4 replace χ by χ_u , we need one injective $\chi_u \in S_A^S$ for each string $u \in \mathcal{L}'_i, i \geq 0$. One mapping χ clearly cannot in general achieve the goal. This small modification changes nothing in the proof of Lemma 3.6.

References

- [1] J. Gustedt, Well-quasi-ordering finite posets, in: N. Robertson and P. Seymour, eds., Proc. AMS-IMS-SIAM Joint Summer Research Conf., Seattle, WA, 1991.
- [2] J. Gustedt, Algorithmic aspects of ordered structures, Ph.D. Thesis, Technische Universität Berlin, 1992.
- [3] G. Higman, Ordering by divisibility in abstract algebras, Proc. London Math. Soc. 2 (1952) 326–336.
- [4] E.C. Milner, Basic wqo and bqo theory, in: I. Rival, ed., Graphs and Orders (Reidel, Dordrecht, 1985).
- [5] R.H. Möhring and R. Müller, A combinatorial approach to obtain bounds for stochastic project networks, Tech. Report, Tech. Univ. Berlin, 1991.
- [6] I. Rival, ed., Graphs and Orders (D. Reidel, Dordrecht, 1985).
- [7] N. Robertson and P. Seymour, eds., Graph minors, Proc. AMS-IMS-SIAM Joint Summer Research Conf., Seattle, WA, 1991.